

Pesquisas em Geociências

<http://seer.ufrgs.br/PesquisasemGeociencias>

New Upper Carboniferous Palaeodictyopteran Insect From Piedra Shotle Formation, Argentina

Irajá Damiani Pinto

Pesquisas em Geociências, 19 (1): 55-58, Mai./Ago., 1992.

Versão online disponível em:

<http://seer.ufrgs.br/PesquisasemGeociencias/article/view/21322>

Publicado por

Instituto de Geociências



Portal de Periódicos UFRGS

UNIVERSIDADE FEDERAL
DO RIO GRANDE DO SUL

Informações Adicionais

Email: pesquisas@ufrgs.br

Políticas: <http://seer.ufrgs.br/PesquisasemGeociencias/about/editorialPolicies#openAccessPolicy>

Submissão: <http://seer.ufrgs.br/PesquisasemGeociencias/about/submissions#onlineSubmissions>

Diretrizes: <http://seer.ufrgs.br/PesquisasemGeociencias/about/submissions#authorGuidelines>

Data de publicação - Mai./Ago., 1992.

Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brasil

New Upper Carboniferous Palaeodictyopteran Insect From Piedra Shotle Formation, Argentina

IRAJÁ DAMIANI PINTO

Instituto de Geociências, UFRGS, Caixa Postal 15001, CEP 91501-970, Porto Alegre, RS, Brasil

(Recebido em 15/04/92. Aceito para publicação em 07/08/92.)

Abstract — A new Palaeodictyopteran insect *Breyeria brauckmanni* from Piedra Shotle Formation, Argentina is described. It belongs to the family Breyeriidae that is attributed to the Carboniferous in North America, Europe and Asia. This fact in connection with the age of the entomofauna and arachnofauna of the Bajo de Veliz Formation that overlies Piedra Shotle Formation, which also is Carboniferous in age, permit to attribute an Upper Carboniferous one for this formation in discordance to the Lower Permian age given to it by Argentinean Palaeobotanists.

Resumo — É descrito um novo inseto da Ordem Palaeodictyoptera, *Breyeria brauckmanni*, proveniente da Formação Piedra Shotle, Argentina. Pertence à Família Breyeriidae, de idade carbonífera nos Estados Unidos da América do Norte, Europa e Ásia. Este fato associado à idade carbonífera da entomofauna e arachnofauna da Formação Bajo de Veliz, sobreposta à Formação Piedra Shotle, permite atribuir a esta formação idade Carbonífero Superior, em discordância com a idade Permiano Inferior, atribuída por paleobotânicos argentinos.

INTRODUCTION

The material under study was kindly provided by Professor Sergio Archangelsky. This material is very important because the presence of insect wings associated with plants. These insects are of exclusively Carboniferous age in several European countries, in the United States of America and in Asia; and on the other hand, the associated taphoflora has a Lower Permian age attributed by some Argentinean Palaeobotanists.

Repository — Museo de Ciencias Naturales, Sector de Paleobotanica, one slab n° BA Pb 637.

GEOGRAPHICAL AND STRATIGRAPHICAL DATA

The insect wing imprint comes from the Bed A of the Profile NFC, La Casilda Section, located at the left margin of the Arroyo Genoa, 3 km North of Piedra Shotle, Chubut, Argentina. The Bed A is 40 cm of a fine laminated light olive grey sediment limited below and above by a coal bed. It contains abundant plant remains which are: *Paranocladus*, *Eucerospermum*, *Aphlebia*, *Pecopteris*, *Sphenopteris*, *Phyllothea*, fructifications and other indeterminated plant remains. The predominant genus is *Paranocladus* with 41.70% frequency. The age is "provisorily Lower Permian". (Archangelsky, 1981). From the coal bed below the bed A were collected the following palynomorphs, according to Gamero and Archangelsky, 1981: *Punctatisporites gretensis* Balme & Hennely, *Lundbladispora braziliensis* (Pant & Srivastava), *Cristatisporites* cf. *lestai* Archangelsky & Gamero. The dominant spores are: *Cristatisporites* and *Lundbladispora* associated with rare remains of *Gangamopteris*. The age attributed for this material is Permian.

TAXONOMY

Ordo Palaeodictyoptera

Familia Breyeriidae Handlirsch, 1906

Sub Familia Breyeriinae Laurentiaux & Laurentiaux-Vieira, 1951

Breyeria de Borre, 1875

Wings of large size, broad, equal in length, the hind pair broader, Sc short or long, stems of R and M approaching or touching near the bases. Branches of main veins few, widely separated from each other; MP and CuA simple, MP forked, CuP usually forked. Hind wing often broadly triangular in shape. Cross-veins numerous, thin, irregular, often connected by anastomoses. Clusters of long hairs on the anterior and posterior margins of the membrane are seen in some species.

Type species: *Pachytylopsis borinensis* de Borre, 1875.

Breyeria brauckmanni Pinto sp. nov.

Figures 1 and 2

Designatio nominis: in honour to Dr Carsten Brauckmann.

Holotypus: one hind wing imprint, Mus. Ci. Nat. n° BA, Pb 637.

Locus typicus: La Casilda Section, Profile NFC, Bed A, Chubut, Argentina.


Diagnosis — Triangular hind wing, with more than 80.0 mm long, 40.0 mm wide; Sc very long fusing with R at almost the apex of the wing; RS with four branches the proximal one forked, the other three single.

Description — Triangular hind wing with the

preserved part 80.0 mm long and 40.0 mm wide. Anterior margin slightly convex, Sc vein parallel to C one and separated from it by straight cross-veins; R runs parallel and close to Sc, separated from it by straight cross-veins also; Rs originated at the base of R runs almost straight, curving backward at the apex and leaving a broader space between it and R specially at the midlength of the wing. This space is covered by anastomosed failbe cross-veins forming a network. Rs has four branches,

the proximal one is bifurcated, MA is single, MP with two long branches, CuA apparently single. Irregular failbe cross-veins cover all the space between Rs, M and CuA.

Remarks — The new species can be differentiated from all other species of the family, except from *B. borinensis* Handlirsch, 1906 and *B. guttata* (Zalessky, 1931), by the long Sc which reaches almost the apex of the wing. Laurentiaux & Laurentiax-Vieira (1951,



Figures 1 and 2 — *Breyeria brauckmanni* Pinto sp. nov. Holotypus Museo de Ciencias Naturales, Buenos Aires, Argentina n° BA Pb 637. Piedra Shotle Formation, Upper Carboniferous, Argentina.

p. 587), say that the drawing of *B. borinensis* (in Handlirsch, 1904, pl. III figs. 9 and 10) presents a Sc vein longer than it is really. However looking the Handlirsch's photograph at the same plate, the difference is very small. Anyway the new species has a longer Sc. The most similar species are: *Tchirkovaea guttata* Zalesky, 1931. [= *Breyeria guttata* (Zalesky, 1931)] and *Borrea lachlani* Brongniart called *Breyeria* (*Borrea*) *lachmani* Brongniart, 1893 (in Laurentiaux & Laurentiaux-Vieira, 1951, p. 590 fig. 4) and *Breyeria lachmani* (Brongniart) (in Kukalová, 1969, Part II, p. 467 fig. 40) which according to the representation made by the author has the same kind of veins disposition; but the new species differs from it in having Sc longer and Rs with one branch more. *Breyeria guttata* (represented in Sinitshenkova, 1979, p. 194 fig. 1) is also greatly similar specially on the long Sc vein but differs in the larger distal separation of Rs from R and by having the posterior branch of the proximal one of Rs (Mp in Sinitshenkova) forked and apparently much stronger cross-veins between Rs, MA, MP and CuA veins.

CONCLUSIONS

The age reached by the taphofauna does not agree with that one attributed to the taphoflora by the following reasons.

a) The world occurrence of the species of the family Breyeriidae is:

Breyeria borinensis (de Borre, 1875), Westphalian C, Belgium.

B. lachlani (Brongniart, 1893), Stephanian B, France.

B. boulei (Meunier, 1910), Stephanian B, France.

B. guttata (Zalesky, 1931), Upper Carboniferous, Siberia.

B. delruei Laurentiaux, 1949, Westphalian B, France.

B. limburgica Laurentiaux, 1950, Westphalian A, Netherlands.

B. stopai Laurentiaux & Laurentiaux-Vieira, 1951, Westphalian C, France.

B. britannica Laurentiaux & Laurentiaux-Vieira, 1951, Westphalian B, England.

B. barborae Kukalová, 1959, Namurian C, Czechoslovakia.

B. vrankeni Laurentiaux-Vieira & Laurentiaux, 1964, Westphalian B, Netherlands.

B. rappi Carpenter, 1967, Westphalian C, USA.

? *B. sharovi* (Sinitshenkova, 1979), Stephanian, Siberia.

? *B. triramosa* (Sinitshenkova, 1979), Stephanian, Siberia.

Stobbsia woodwardiana Handlirsch, 1908, Westphalian, England.

[= *Breyeria woodwardiana* (Handlirsch, 1906) in Bolton, 1921].

Breyerioides kliveri (Güthörl, 1934), Westphalian C, Germany.

Jugobreyeria sippelorum Brauckmann, 1985, Namurian B, Germany.

So the insect under description belongs to a family until now being exclusively of Carboniferous age.

b) Another insect under description from Piedra Shotle Formation belongs also to a group of exclusively Carboniferous occurrence.

c) Bajo de Veliz Formation overlying the Piedra Shotle Formation has a taphofauna (insects and arachnids) of exclusively Carboniferous age (Pinto & Ornellas, 1978a, b; 1980a, b; 1991; Pinto & Hünicken, 1980; Pinto, Ornellas & Purper, 1980).

Based on these facts the author concludes an Upper Carboniferous age for the Piedra Shotle Formation.

Acknowledgments — Many thanks to Professor Sergio Archangelsky who kindly provided the material for study; to Professor Cândido Simões Ferreira for the help in bibliography; to Lizeth Cardoso Marques for helping in the drawing. To Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and to Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS) for the author's researches continuous support.

REFERENCES

- Archangelsky, S. 1981. Estudios fitopaleoecológicos en el Paleozoico Superior del Oeste de Chubut, Argentina. *Anais II Congresso Latino-Americano de Paleontologia*, Porto Alegre, 1: 141-156.
- Bolton, H. 1921. A Monograph of the Fossil Insect of the British Coal Measures. *Paleontographical Society Monograph*, 74: 1-80.
- Borre, A.P. de 1875a. Note sur des empreintes fossiles decouvertes dans les schistes houillers des environs de Mons. *Compte Rendu Société Entomologique Belge*, 12 (2): 4-7.
- 1875b. Complement de la note sur des empreintes des insectes fossiles. *Compte Rendu Société Entomologique Belge*, 18 (2): 7-11.
- Brauckmann, C.; Koch, L. & Kemper, M. 1985. Spinnentiere (Arachnida) und Insekten aus den Vorhaller-Schichten (Namurium B. Ober-Karbon) Von Hagen-Vorhalle (West-Deutschland). *Geologie und Paläontologie in Westfalen*, 3: 1-132.
- Brongniart, C. 1893. Recherches pour servir à l'Histoire des Insectes Fossiles des Temps Primaires précédées d'une étude sur la nervation des ailes des insectes. Saint Etienne, Theolier, 493p.
- 1893. Recherches pour servir à l'Histoire des Insectes Fossiles des Temps Primaires précédées d'une étude sur la nervation des ailes des insectes: atlas. Saint Etienne, Theolier, 43p.
- Carpenter, F.M. 1944. Carboniferous Insects from the Vicinity of Mazon Creek, Illinois. *Illinois State Museum, Science Papers*, 3 (1): 1-20.
- Gamero, J.C. & Archangelsky, S. 1981. Halazgo de Palinomorfos Pérmicos en Sedimentitas de la Formación Piedra Shotle, Estancia La Casilda y en la Perforación Canadón, Pastos Blancos (YPF Ch es-1), Chubut, Argentina. *Anais, II Congresso Latino-Americano de Paleontologia*, Porto Alegre, 1: 169-177.
- Güthörl, P. 1934. Die Arthropoden aus dem Carbon und Perm des Saas-Nahe-Pflaz-Gebietes. *Abhandlungen der Preussischen Geologischen Landesanstalt*, 164: 1-219.
- Handlirsch, A. 1904. Les Insectes Houillers de la Belgique. *Memoires du Musée Royal d'Histoire Naturelle de Belgique*, 3: 1-20.
- 1906. Revision of American Paleozoic Insects. *Proceedings of the United States National Museum*, 29 (1441): 661-820.
- 1906-1908. Die fossilen Insekten und die Phylogenie des rezenten Formen. *Handbuch für Paleontologie Engelmann*, 1-6: 1-1433.
- Haupt, H. 1949. Rekonstruktionen permokarbonischer Insekten. *Beiträge zur Taxonomischen Zoologie*, 1: 23-43.
- Kukalová, J. 1959. *Breyeria barborae* n. sp. (Insecta, Palaeodictyoptera) of Upper Silesian coal basin (Westphalian). *Vestník Ústředního Ústavu Geologického*, 34: 310-313.
- 1969a. Revisional study of the order Palaeodictyoptera in the Upper Carboniferous Shales of Commentry, France. Part 1. *Psyche*, 76 (2): 163-215.

- . 1969b. Revisional study of the order Palaeodictyoptera in the Upper Carboniferous Shales of Comentry, France. Part II. *Psyche*, 76 (4): 439-468.
- Laurentiaux, D. 1949. Un insecte nouveau Breyeriidae du terrain houiller du Pas-de-Calais. *Annales Société Géologique du Nord*, 99: 47-54.
- . 1950. Les Insectes Houillers du Limbourg Hollandais. *Mededelingen van den Geologische Stichting. N. Ser.*, 4: 13-23.
- & Laurentiaux-Vieira, F. 1951. Observations sur les Paléodictyoptères Breyeriidae. *Bulletin de la Société Géologique de France, 1er. Ser.*, 66 (7): 585-596.
- & ———. 1963. Nouvelle Recolte d'un Breyeridae (Insecte Paléodictyoptère) dans le Westphalien du Nord de la France. *Annales Société Géologique de Nord*, 82: 173-178.
- & ———. 1964. Diagramme pteronologique du genre Westphalien Breyeria de Borre (Insectes, Paléodictyoptère). *Compte Rendu Académie Science*, 238 (9): 1282-1284.
- Meunier, F. 1908. Un Platypteride et un Sténodictyopteride du Stephanien de Comentry. *Annales Société Science*, 32 (2): 153-154.
- . 1908. Nouveau Paléodictyoptère du Stephanien de Comentry. *Bulletin du Muséum d'Histoire Naturelle*, 14: 34-36.
- . 1909. Nouvelles recherches sur les insectes du Terrain Houiller de Comentry (Allier). *Annales de Paléontologie*, 4: 125-152.
- . 1910. Nouveaux Paléodictyoptères du Terrain Houiller de Comentry. *Bulletin du Muséum d'Histoire Naturelle*, 4: 233-237.
- . 1912. Nouvelles recherches sur quelques insectes du Terrain Houiller de Comentry (Allier). Part 2. *Annales de Paléontologie*, 7: 3-19.
- Pinto, I.D. & Hünicken, M.A. 1980. Gondwanarachne a new genus of the Order Trigonotarbitida (Arachnida) from Argentina. *Boletín de la Academia Nacional de Ciencias*, 53 (3-4): 307-315.
- & Ornellas, L.P. de 1978a. Upper Carboniferous Insects from Argentina. I. Familia Diaphanopteridae (Megasecopteroidea). *Pesquisas*, 10: 87-95.
- & ———. 1978b. Carboniferous Insects (Protorthoptera and Paraplecoptera) from the Gondwana (South America, Africa and Asia). *Pesquisas*, 11: 305-321.
- & ———. 1980a. Upper Carboniferous Insects from Argentina. II. Narkemocacurgidae (Paraplecoptera). *Boletín de la Academia Nacional de Ciencias*, 53 (3-4): 287-291.
- & ———. 1980b. A New Upper Carboniferous Paraplecopteran Insect from Argentina. *Anais, II Congresso Latino-Americano de Paleontologia*, Porto Alegre, 1: 108-110.
- & ———. 1991. Substitute Names for the Extinct Insecta Families Narkemocacurgidae Pinto & Ornellas, 1978 and Cacurgonarkemidae Pinto, 1990. *Pesquisas*, 18 (1): 93.
- ; ——— & Purper, I. 1980. Fossil insects and the problem of Permo-Carboniferous Boundaries. *Actas, Segundo Congreso Argentino de Paleontología y Bioestratigrafía y Primer Congreso Latinoamericano de Paleontología*, Buenos Aires, Asociación Paleontologica Argentina, 4: 207-213.
- Pruvost, P. 1930. La faune continentale du terrain houiller de la Belgique. *Mémoire du Muséum Royal d'Histoire Naturelle de Belgique*, 44: 1-146.
- Rodendorf, B.B. 1962. *Osnovy paleontologii-Arthropoda, Tracheata and Chelicerata*. Izdatelstvo Akademii Nauk SSSR. 560 p.
- Schneider, J. 1983. Taxonomie und Palaökologie der Blattodea-Fauna aus dem Stefan von Comentry (Frankreich). Versuch einer Revision. *Freiberg Forschungsheft*, C376: 77-98.
- Sinitshenkova, N.D. 1979. A new family of the Palaeodictyoptera from the Carboniferous of Siberia. *Paleontological Journal*, 13 (2): 192-205.
- Zalessky, M. 1931. On insects found at the coal deposits of the Kuznetsk basin and the age based on the entomofauna. *Biologie Moskaw Prirody*, 3/4: 399-406. (In Russian).